Osgood-Schlatter's Disease

IF YOU NOTICE PAIN JUST BELOW THE KNEE, IT COULD BE OSGOOD-SCHLATTER'S.

NOT TO WORRY, THOUGH, IT WILL SUBSIDE AND DRASTIC TREATMENT IS SELDOM NECESSARY.

By Andrew DeGruccio, MD

sgood-Schlatter's disease is always characterized by activity-related pain that occurs a few inches below the knee-cap, or patella, on the front of the knee. The child will have swelling in the area, and tenderness to the touch. Sports requiring lots of running, jumping, kneeling and squatting are particularly associated with this disorder. Many children first signal the start of the problem by rubbing the top of their "shinbones" with their hands, or even ice cubes, at practice sessions. The three main factors that contribute to Osgood-Schlatter's are:

- 1. Between 10 and 15 years old
- 2. Involved in youth sports
- 3. In a "growth spurt"

The condition is caused by the powerful quadriceps muscle pulling on the attachment point of the patellar tendon during running activities such as soccer, basketball, track and other sports. The patellar tendon is a thick rope-like structure that connects the bottom of the kneecap (patella) to the top of the large shin bone (tibia). The powerful muscles on the front of the thigh, the quadriceps muscles, straighten the knee by pulling at the patellar tendon via the patella. Osgood-Schlatter's disease is caused by inflammation (irritation) where the patellar tendon attaches to the tibia. The disease is caused by repetitive stress or tension on part of the growth area of the upper tibia (the apophysis). The disease may also be associated with an injury in which the tendon is stretched so much that it tears away from the tibia and takes a fragment of bone with it.

If you have Osgood-Schlatter's, sport or physical activity is not likely to cause any permanent damage, but may make the pain worse. Difficult decisions may have to be made about training schedules and sporting commitments. Each case is different and your doctor will advise. An approach similar to the following may be advised.

You do not usually need to stop sport. However, easing off strenuous or vigorous sport may be sensible for a while until the pain eases to a tolerable level. Let the pain be your guide. Play sport only to a bearable level of pain. Avoid heavy sport activities if the pain is bad. Consider a complete break from sport for a while if the pain remains bad.

An ice pack applied below the kneecap for about 5 minutes before and after exercise may prevent some inflammation and pain.

Consider wearing knee pads during sports such as football, volleyball, and basketball to protect the tender area.

Concentrate on flexibility and quadriceps stretching, especially during growth spurts.

Anti-inflammatory medications, i.e. ibuprofen, are often very helpful.

Prevention of the problem has been low on the priority list for most healthcare providers. This may be because prevention techniques are unproven and controversial. Better pre-sport conditioning of the quadriceps may help prevent the problem entirely. Clearly, better flexibility and stretching throughout adolescence may be crucial.

Osgood-Schlatter's disease is not "torn ligaments" or "water on the knee," nor is it the end of athletics, but many sources of information out there can be misleading and confusing. Entire websites have been dedicated to this diagnosis, some promoting herbal supplements, others requesting your money to provide the very information that was just discussed. Some sites out of Europe even discuss surgical options for this condition. Rest assured, surgery is not the recommended modality of treatment for this in the United States, nor is prolonged immobilization.

In essence, it is a self-limited adolescent condition that is fairly common in athletic kids, with no real permanent disabling sequelae except for a bump just below the knee. The pain from the condition usually subsides by the time the athlete is 15 to 16 years old, and only rarely are there any residual symptoms associated with the visible bump.•

